arXiv:1503.02547v3 [math.GT] 20 Apr 2015

A volume conjecture for a family of Turaev-Viro type invariants of 3-manifolds with boundary

Qingtao Chen & Tian Yang

Abstract

We consider a family of Turaev-Viro type invariants for a 3-manifold M with non-empty boundary, indexed by an integer $r \geqslant 3$, and propose a volume conjecture for hyperbolic M that these invariants grow exponentially at large r with a growth rate the hyperbolic volume of M. The crucial step is the evaluation at the root of unity $\exp(2\pi\sqrt{-1}/r)$ instead of that at the usually considered root $\exp(\pi\sqrt{-1}/r)$. Evaluating at the same root $\exp(2\pi\sqrt{-1}/r)$, we then conjecture that, the Turaev-Viro invariants and the Reshetikhin-Turaev-Lickorish invariants of a closed hyperbolic 3-manifold M grow exponentially with growth rates respectively the hyperbolic and the complex volume of M. This uncovers a different asymptotic behavior of the values at other roots of unity than that at $\exp(\pi\sqrt{-1}/r)$ predicted by Witten's Asymptotic Expansion Conjecture, which may indicate some different geometric interpretation of the Reshetikhin-Turaev invariants than the $SU(2)$ Chern-Simons theory. Numerical evidences are provided to support these conjectures.

1 Introduction

In [24, 25], Kashaev introduced a family of complex valued link invariants indexed by the integers n using quantum dilogarithm, and conjectured that, for a hyperbolic knot or link, the absolute value of these invariants grow exponentially at large n, with a growth rate the hyperbolic volume of the knot or link complement in S^3. The mathematically rigorous definition of Kashaev's invariants was given by Baseilhac and Benedetti [1, 2, 3, 4]. In [39], Murakami and Murakami proved that Kashaev's invariants turned out to be special values of the colored Jones polynomials of the link, and extend Kashaev's conjecture to the following

Volume Conjecture ([25, 39]). For any link K in S^3, let $J_n(K;q)$ be its n-th colored Jones polynomial. Then

$$\lim_{n\to+\infty}\frac{2\pi}{n}\ln\left|J_n(K;e^{\frac{\pi\sqrt{-1}}{n}})\right| = v_3\cdot\|S^3\setminus K\|,$$

where v_3 is the volume of the regular hyperbolic ideal tetrahedron and $\|S^3\setminus K\|$ is the simplicial volume of the complement of K.

Due to the use of different notations, the value $J_n(K;e^{\frac{\pi\sqrt{-1}}{r}})$ here equals the value of the colored Jones polynomial at the usually considered root of unity $e^{\frac{2\pi\sqrt{-1}}{r}}$ in most of the other literatures. (See also Remark 2.1.) This conjecture has been proved for certain cases including the Figure-eight knot (cf. [39]), torus knots [26] and their Whitehead doubles [57], $(2,2m)$-torus links [23], Borromean rings [18], twisted Whitehead links [57] and Whitehead chains [54]. In [55], it was proved that for every link K there is an unlink L in the complement of K such that the Volume Conjecture is true for the union $K\cup L$. Numerically, the Volume Conjecture has been verified for the knots 6_3, 8_9 and 8_{20} [40]. Various extensions of this conjecture for the colored Jones polynomials were proposed and proved for certain cases in [22, 41, 10, 15, 14, 42].

In the present paper, we consider a family of real valued invariants $\{TV_r(M;q)\}$ for a compact 3-manifold M with non-empty boundary, indexed by an integer $r\geqslant 3$ and evaluated at certain root of unity q of degree $2r$, which is an extension of Turaev-Viro invariants for closed 3-manifolds [52]. (See Definition 3.1 and Theorem 3.2.) Then main focus of this work is on the valus at the root of unit $q=e^{\frac{2\pi\sqrt{-1}}{r}}$ and their asymptotic behavior, and we propose the following volume conjecture.

Conjecture 1.1. *Let M be a hyperbolic 3-manifold with cusps or geodesic boundary and let $Vol(M)$ be its hyperbolic volume. Then for r running over all odd integers and for $q = e^{\frac{2\pi\sqrt{-1}}{r}}$,*

$$\lim_{r\to+\infty}\frac{2\pi}{r-2}\log\big(TV_r(M;q)\big) = Vol(M).$$

The invariant $TV_r(M)$ is a state sum of Turaev-Viro type defined by using triangulations and quantum $6j$-symbols (§2.2). Recall that Turaev-Viro's original construction gives rise to real valued invariants for closed 3-manifolds and $2{+}1$ TQFT's for 3-cobordisms [52], and $2{+}1$ TQFT's for 3-cobordisms containing a link inside [50]. In all of their constructions, they use the usual triangulations, meaning the vertices of the triangulations are inside the manifolds and the cobordisms, and in the case of cobordisms, the edges on the boundary are from the edges of the triangulations. The difference in our construction is that, instead of using the usual triangulations, we use *ideal triangulations* of a 3-manifold with non-empty boundary (§2.1). Then following Turaev-Viro's construction, we color the edges by half-integers less than or equal to $(r-2)/2$, associating a quantum $6j$-symbol to each tetrahedron. We then multiply the quantum $6j$-symbols over all the tetrahedra and sum up the resulting products over all the colorings to obtain a number. The Orthogonality and Biedenharn-Elliot identity assure that this real number is independent of the choice of the ideal triangulation, and hence is an invariant of the 3-manifold. See §3 for more details. For orientated 3-manifolds with non-empty boundary, we are informed by Benedetti that these invariants coincide with the ones defined in [5] using o-graphs.

Conjecture 1.1 is motivated by the beautiful work of Costantino [11] that the quantum $6j$-symbols for a sequence of 6-tuples of hyperbolic type grow exponentially, with the growth rate the hyperbolic volume of the truncated hyperideal tetrahedron whose dihedral angles are determined by the limit of the 6-tuples. See also [13, 14]. There are plenty of supporting evidences to Conjecture 1.1. In §4, we numerically calculate $TV_r(M)$ for the Figure-eight knot complement and its sister, for the knot 5_2 complement and the manifolds M_{3_6} and M_{3_8} in Callahan-Hildebrand-Weeks' census [8]. Recall that M_{3_6} is the complement of a 12-crossings knot and has the same volume as the knot 5_2 complement. We also numerically calculate $TV_r(M)$ for the knot 6_1 complement, for the non-orientable cusped 3-manifolds N_{1_1} and N_{2_1} in Callahan-Hildebrand-Weeks' census, and for the smallest hyperbolic 3-manifolds with a totally geodesic boundary [17, 34]. The results turn out to be very promising.

The invariants described above can further be defined as invariants for closed pseudo-3-manifolds and, in the case that the underling space of the pseudo-manifold is a closed 3-manifold, differ from the original Turaev-Viro invariant by a constant depending on and growing polynomially in r. Based on this and Conjecture 1.1, we propose the following volume conjecture for the original Turaev-Viro invariants [52] for closed 3-manifolds.

Conjecture 1.2. *Let M be a closed hyperbolic 3-manifold, let $TV_r(M;q)$ be its Turaev-Viro invariant and let $Vol(M)$ be its hyperbolic volume. Then for r running over all odd integers and for $q = e^{\frac{2\pi\sqrt{-1}}{r}}$,*

$$\lim_{r\to+\infty}\frac{2\pi}{r-2}\log\big(TV_r(M;q)\big) = Vol(M).$$

Evidences to Conjecture 1.2 are provided in §5. The calculation is based on a relationship between the Turaev-Viro invariants and the Reshetikhin-Turaev-Lickorish invariants [45, 46, 31, 47]. Recall that the Reshetikhin-Turaev invariants $\{\tau_r(M;q)\}$ are complex valued invariants of oriented closed 3-manifolds, defined for all integers $r \geqslant 3$ and for all the primitive roots of unity q of degree $2r$ using representation theory of quantum groups. For $q = e^{\frac{\pi\sqrt{-1}}{r}}$, these invariants provided a mathematical construction of the 3-manifold invariants introduced by Witten [53] using quantum field theory with Chern-Simons action. In [29, 30, 31], Lickorish defined Reshetikhin-Turaev invariants using skein theory and extended $\tau_r(M;q)$ for odd r to the primitive roots of unit q of degree r. In particular, when r is odd, $\tau_r(M;q)$ was defined at $q = e^{\frac{2\pi\sqrt{-1}}{r}}$. Numerically calculating $\tau_r(M;q)$ for closed 3-manifolds obtained from S^3 by doing certain integral Dehn-surgeries along the Figure-eight knot and the 5_2 knot in §5, we propose the following

Conjecture 1.3. *Let M be a closed oriented hyperbolic 3-manifold and let $\tau_r(M;q)$ be its Reshetikhin-Turaev-Lickorish invariants. Then for r running over all odd integers and $q = e^{\frac{2\pi\sqrt{-1}}{r}}$, and with a suitable choice of the arguments,*

$$\lim_{r\to+\infty}\frac{4\pi}{r-2}\log\big(\tau_r(M;q)\big) \equiv Vol(M) + \sqrt{-1}CS(M) \quad (mod\ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

where $CS(M)$ is the usual Chern-Simons invariants of M multiplied by $2\pi^2$.

Conjecture 1.3 is quite unexpected since it is previously widely believed that only the value of $\tau_r(M;q)$ at $q = e^{\frac{\pi\sqrt{-1}}{r}}$ is pivotal and the values at the other roots of unity can be obtained from the former by certain Galois transformations. Compared with Witten's Asymptotic Expansion Conjecture (cf. [43]), we can see that values of $\tau_r(M;q)$ at other roots of unity, in particular at $q = e^{\frac{2\pi\sqrt{-1}}{r}}$, have a rather different asymptotic behavior, and we expect a geometric and physical interpretation of this new phenomenon.

One of the significances of Kashaev-Murakami-Murakami Volume Conjecture is that it relates the quantum invariants and the hyperbolic geometry. In particular, it provides a possible method to calculate the hyperbolic volume. The traditional method of calculating the volume is either to solve Thurston's hyperbolic gluing equations and apply the dilogarithm functions [48], or to solve Casson's angle structure equations and apply the volume optimization process [9]. Both of these approaches work for cusped hyperbolic 3-manifolds only. In [32], Luo generalized Casson's program to hyperbolic 3-manifolds with totally geodesic boundary. However, all these methods heavily rely on the ideal triangulations. For a bad choice of the triangulation, the solutions to both systems of equations may not exist. Assuming the Volume Conjecture, the simplicial volume of a knot or link complement could be calculated as the growth rate of values of the colored Jones polynomials. However, this method will only work for knot or link complements in S^3. In [1, 2, 3, 4], Kashaev's invariants were generalized by Baseilhac-Benedetti to link complements in any closed 3-manifolds, and hence could potentially be used to calculate the volume there. If Conjectures 1.1, 1.2 and 1.3 are proved true, then they will provide a new approach to calculate the hyperbolic and the complex volume, i.e., as the growth rates of $TV_r(M)$ and $\tau_r(M)$. This method will work for all compact 3-manifolds either with or without boundary, and will work for any triangulations. Numerical evidences also suggest that, for the knot and link complements, it is more efficient than using the colored Jones polynomials. For non-hyperbolic 3-manifolds that are closed or with torus boundary, the above conjectures can be asked for the convergence to and be used to calculate the simplicial volumes as the Kashaev-Murakami-Murakami Volume Conjecture.

In §6, we calculate $TV_r(M)$ for the complements of the unknot U, the Hopf link, the Trefoil knot and the torus links $T_{(2,4)}$ and $T_{(2,6)}$. In particular, we show that $TV_r(S^3 \setminus U) = 1$ for all $r \geqslant 3$ and for all possible roots of unity q (Proposition 6.1), and we conjecture that the family $\{TV_r(M)\}$ detects the unknot (Conjecture 6.2). Numerically, we also calculate $TV_r(M)$ for the torus knots $T_{(2,5)}$, $T_{(2,7)}$, $T_{(2,9)}$, $T_{(2,11)}$, $T_{(3,5)}$ and $T_{(3,7)}$ complements. All the calculations above suggest the following integrality conjecture.

Conjecture 1.4. *Let $T_{(m,n)}$ be the (m,n)-torus link in S^3, and suppose $m < n$. If r is coprime to n, then $TV_r(S^3 \setminus T_{(m,n)})$ is an integer independent of the choice of the roots of unity q.*

The unexpected emergence of integers in Conjecture 1.4 suggests that there might be some nontrivial identities for the quantum $6j$-symbols behind, and we hope this phenomenon will shed lights to the representation theory of quantum groups. In §7, we compare the asymptotic behavior of $TV_r(M)$ for a link complement to that of the values of the corresponding colored Jones polynomials, and propose some questions for the further studies.

Acknowledgments: The authors are grateful to Reccardo Benedetti, Francis Bonahon and Nicolai Reshetikhin for inspiring discussions and suggestions and Hao Zheng for helpful conversations. The first author would like to thank Kefeng Liu and Weiping Zhang for a lot of useful discussions during the past few years. The second author is grateful to Henry Segerman and Hongbin Sun for helpful discussions and Steven Kerckhoff, Feng Luo and Maryam Mirzakhani for suggestions.

The first author is supported by a grant associated to his Senior Shiing-Shen Chern Postdoctoral Fellowship from ICTP. The second author is supported by NSF grant DMS-1405066.

2 Preliminaries

2.1 Ideal triangulations

Ideal triangulations of a 3-manifold with boundary were first introduced by Thurston in [48] and used to find the hyperbolic structure on the Figure-eight knot complement. After Thurston, ideal triangulations became a main tool of doing various calculations on 3-manifolds. In our construction, the use of ideal triangulations is also one of the key ingredients.

Let M be a compact 3-manifold with non-empty boundary. An *ideal triangulation* $\mathcal{T}$ of M consists of a disjoint union $X = \bigsqcup \Delta_i$ of finitely many oriented truncated Euclidean tetrahedra Δ_i and a collection of orientation reversing homeomorphisms Φ between pairs of hexagonal faces in X. The quotient space X/Φ is a compact 3-manifold M with a triangulation $\mathcal{T}$ where the boundary of M is the quotient of the triangular faces in X. An edge of a truncated Euclidean tetrahedra is the intersection of two hexagonal faces. The *edges* in $\mathcal{T}$ are the quotients of the edges in X, the *faces* in $\mathcal{T}$ are the quotient of the hexagonal faces in X, and the *tetrahedra* in $\mathcal{T}$ are the quotients of the truncated Euclidean tetrahedra in X. In [36, 37, 44], it is proved that any two ideal triangulations of M are related by a sequence of $0-2$ and $2-3$ Pachner moves. See the figure below.



0-2 move



2-3 move

2.2 Quantum $6j$-symbols

The other key ingredient in our construction is the quantum $6j$-symbols associated with the quantized universal enveloping algebra $U_q(sl_2(\mathbb{C}))$ developed in [28]. In this subsection, we will recall the definition and some basic properties we need, and we refer to [28, 27] for more details about the quantum $6j$-symbols.

Through out this subsection, we will fix an integer $r \geqslant 3$, and let $I_r = \{0, 1/2, \dots, (r-2)/2\}$ be the set of half-integers less than or equal to $(r-2)/2$. Let q be a root of unity of degree $2r$ such that q^2 is a primitive root of unity of degree r. For an integer n, the quantum number $[n]$ is the real number defined by

$$[n] = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

In particular, when $q = e^{k\pi\sqrt{-1}/r}$,

$$[n] = \frac{\sin(nk\pi/r)}{\sin(k\pi/r)}.$$

The quantum factorial is defined by $[n]! = [n][n-1]\dots[1]$, and as a convention, $[0]! = 1$.

Remark 2.1. The quantum number we use in this paper is different from some of the other literatures where in their notation $[n] = (q^{n/2} - q^{-n/2})/(q^{1/2} - q^{-1/2})$.

A triple (i, j, k) of elements of I_r is called *admissible* if (i) $i + j \geqslant k$, $j + k \geqslant i$ and $k + i \geqslant j$, (2) $i + j + k$ is an integer, and (3) $i + j + k \leqslant r - 2$. For an admissible triple (i, j, k), define

$$\Delta(i, j, k) = \sqrt{\frac{[i + j - k]![j + k - i]![k + i - j]!}{[i + j + k + 1]!}}.$$

As a convention here, we let $\sqrt{x} = \sqrt{-x}\sqrt{-1}$ for a negative real number x.

Quantum $6j$-symbols. Let (i, j, k, l, m, n) be a 6-tuple of elements of I_r such that the triples (i, j, k), (j, l, n), (i, m, n) and (k, l, m) are admissible. Let $T_1 = i + j + k$, $T_2 = j + l + n$, $T_3 = i + m + n$ and $T_4 = k + l + m$, and let $Q_1 = i + j + l + m$, $Q_2 = i + k + l + n$ and $Q_3 = j + k + m + n$. Then the quantum $6j$-symbol for the

6-tuple (i,j,k,l,m,n) is defined by

$$\left|\begin{array}{ccc} i & j & k \\ l & m & n \end{array}\right| = \sqrt{-1}^{-2(i+j+k+l+m+n)}\Delta(i,j,k)\Delta(j,l,n)\Delta(i,m,n)\Delta(k,l,m)$$
$$\times \sum_{k=\max\{T_1,T_2,T_3,T_4\}}^{\min\{Q_1,Q_2,Q_3\}} \frac{(-1)^z[z+1]!}{[z-T_1]![z-T_2]![z-T_3]![z-T_4]![Q_1-z]![Q_2-z]![Q_3-z]!}.$$

A good way to memorize this definition is to use a tetrahedron. As in the figure below, each T_i corresponds to a face of the tetrahedron, and each Q_j corresponds to a quadrilateral in the tetrahedron.



From the definition, it is easy to see that

$$\left|\begin{array}{ccc} i & j & k \\ l & m & n \end{array}\right| = \left|\begin{array}{ccc} j & i & k \\ m & l & n \end{array}\right| = \left|\begin{array}{ccc} i & k & j \\ l & n & m \end{array}\right| = \left|\begin{array}{ccc} i & m & n \\ l & j & k \end{array}\right| = \left|\begin{array}{ccc} l & m & k \\ i & j & n \end{array}\right| = \left|\begin{array}{ccc} l & j & n \\ i & m & k \end{array}\right|.$$

The quantum $6j$-symbols satisfy the follow two important identities, which are the key properties for defining our invariants. For $i \in I_r$, define

$$w_i = (-1)^{2i}[2i+1].$$

Orthogonality. For any $i,j,k,l,m,n \in I_r$ such that (i,j,m), (i,j,n), (k,l,m) and (k,l,n) are admissible (see the figure below), we have

$$\sum_s w_s w_m \left|\begin{array}{ccc} i & j & m \\ k & l & s \end{array}\right| \left|\begin{array}{ccc} i & j & n \\ k & l & s \end{array}\right| = \delta_{mn}, \tag{2.1}$$

where δ is the Kronecker symbol, and the sum is over $s \in I_r$ such that the 6-tuples involved in the sum are admissible.



Biedenharn-Elliot identity. For any $i,j,k,l,m,n,o,p,q \in I_r$ such that (o,p,q,i,j,k) and (o,p,q,l,m,n) are admissible (see the figure below), we have

$$\sum_s w_s \left|\begin{array}{ccc} i & j & q \\ m & l & s \end{array}\right| \left|\begin{array}{ccc} j & k & o \\ n & m & s \end{array}\right| \left|\begin{array}{ccc} k & i & p \\ l & n & s \end{array}\right| = \left|\begin{array}{ccc} o & p & q \\ i & j & k \end{array}\right| \left|\begin{array}{ccc} o & p & q \\ l & m & n \end{array}\right|, \tag{2.2}$$

where the sum is over $s \in I_r$ such that the 6-tuple involved in the sum are admissible.



3 Turaev-Viro type invariants on ideal triangulations

In this section, we describe the Turaev-Viro type invariants $TV_r(M)$ of a 3-manifold M with non-empty boundary, using ideal triangulations of M. Note that we do not put any constrains on the orientability nor on the type of the boundary components of M, they could be either tori or high genus surfaces. For orientated 3-manifolds with boundary, $\{TV_r(M)\}$ coincide with the ones defined in [5].

For $r \geqslant 3$ let $I_r = \{0, 1/2, 1, \dots, r-2/2\}$. A *coloring* of a Euclidean tetrahedron Δ is an assignment of elements of I_r to the edges of Δ, and is *admissible* if the triple assigned to the three edges of each face of Δ is admissible. Let $\mathcal{T}$ be an ideal triangulation of M. A *coloring at level* r of the ideally triangulated 3-manifold $(M, \mathcal{T})$ is an assignment of elements of I_r to the edges of $\mathcal{T}$, and is *admissible* if the 6-tuple assigned to the edges of each tetrahedron of $\mathcal{T}$ is admissible.

Let q be a root of unity of degree $2r$ such that q^2 is a primitive root of unity of degree r, and for any integer n, let $[n]$ be the quantum number defined in §2.2. Let c be an admissible coloring of $(M, \mathcal{T})$ at level r. For each edge e of $\mathcal{T}$, we let

$$|e|_c = w_{c(e)}.$$

For each tetrahedra Δ in $\mathcal{T}$ with vertices $v_1, \dots, v_4$, we denote by e_{ij} the edge of Δ connecting the vertices v_i and v_j, $\{i, j\} \subset \{1, \dots, 4\}$, and let

$$|\Delta|_c = \begin{vmatrix} c(e_{12}) & c(e_{13}) & c(e_{23}) \\ c(e_{34}) & c(e_{24}) & c(e_{14}) \end{vmatrix}.$$

Definition 3.1. *Let $A_r(M, \mathcal{T})$ be the set of admissible colorings of $(M, \mathcal{T})$ at level r, and let E and T respectively be the sets of edges and tetrahedra in $\mathcal{T}$. Define*

$$TV_r(M, \mathcal{T}) = \sum_{c \in A_r(M, \mathcal{T})} \prod_{e \in E} |e|_c \prod_{\Delta \in T} |\Delta|_c.$$

Theorem 3.2. *For each r and q as above, $TV_r(M, \mathcal{T})$ is a real number independent of the choice of $\mathcal{T}$, i.e., defines an invariant of M.*

In the rest of the paper, we will omit the triangulation $\mathcal{T}$, and denote the invariant by $TV_r(M)$, and by $TV_r(M; q)$ if the root of unity q needs to be specified.

Proof of Theorem 3.2. From the definition, there are two possibilities for a quantum $6j$-symbol of the 6-tuple (i, j, k, l, m, n) to be a non-real complex number. One is from the square root in $\Delta(i, j, k)$, $\Delta(j, l, n)$, $\Delta(i, m, n)$ and $\Delta(k, l, m)$, and the other is from the coefficient $\sqrt{-1}^{-2(i+j+k+l+m+n)}$. Since each triangle of $\mathcal{T}$ is adjacent to two tetrahedra, the square roots in $\Delta(i, j, k)$'s show up in pairs in $TV_r(M)$, which together give rise to a real number. To the coefficient $\sqrt{-1}^{-2(i+j+k+l+m+n)}$, the triangle of $\mathcal{T}$ with a coloring (i, j, k), say, contributes a factor $\sqrt{-1}^{-(i+j+k)}$. Together with the same factor from the other tetrahedral adjacent to it, this triangle colored by (i, j, k) gives rise to a factor $(-1)^{-(i+j+k)}$ to the summand of $TV_r(M)$. Since the coloring is admissible,

$i+j+k$ is an integer, and hence $(-1)^{-(i+j+k)} = \pm 1$ is a real number. Therefore, each summand in the definition of $TV_r(M)$ is real, which implies that $TV_r(M)$ is real.

By [36, 37, 44], to show that $TV_r(M)$ is independent of the choice of the ideal triangulation, it suffices to prove that $TV_r(M)$ is invariant under the $0-2$ and $2-3$ Pachner moves. For the former case, we suppose that $\mathcal{T}'$ is an ideal triangulation of M obtained from an ideal triangulation $\mathcal{T}$ by doing a $0-2$ Pachner move. Let E and T respectively be the sets of edges and tetrahedral of $\mathcal{T}$, and let E' and T' be the sets of edges and tetrahedra of $\mathcal{T}'$. Suppose e_1 is the edge of $\mathcal{T}$ where the $0-2$ move is performed, and $e_2, e_3 \in E'$ and $\Delta_1', \Delta_2' \in T'$ are respectively the new edges and tetrahedra in $\mathcal{T}'$. Then $E' \setminus \{e_2, e_3\} = E$ and $T' \setminus \{\Delta_1', \Delta_2'\} = T$. See the figure below.



We have

$$\begin{aligned} TV_r(M,\mathcal{T}') &= \sum_{c\in A_r(M,\mathcal{T}')} \prod_{e\in E'} |e|_c \prod_{\Delta\in T'} |\Delta|_c \\ &= \sum_{c\in A_r(M,\mathcal{T}')} \prod_{e\in E'\setminus\{e_2\}} |e|_c \prod_{\Delta\in T'\setminus\{\Delta_1',\Delta_2'\}} |\Delta|_c \Big(\sum_s |e_2|_s |\Delta_1'|_{c,s} |\Delta_2'|_{c,s}\Big) \\ &= \sum_{c\in A_r(M,\mathcal{T}')} \prod_{e\in E'\setminus\{e_1,e_2\}} |e|_c \prod_{\Delta\in T'\setminus\{\Delta_1',\Delta_2'\}} |\Delta|_c \Big(\sum_s |e_2|_s |e_1|_c |\Delta_1'|_{c,s} |\Delta_2'|_{c,s}\Big), \end{aligned} \tag{3.1}$$

where the second sum is over all $s \in I_r$ such that the triples involved are admissible, $|e_2|_s = w_s$, and $|\Delta_1'|_{c,s}$ and $|\Delta_2'|_{c,s}$ are respectively the quantum $6j$-symbols of the 6-tuples assigned by the colorings c and s to the edges of Δ_1' and Δ_2'. By the Orthogonality (2.1), the last term of equation (3.1) equals

$$\sum_{c\in A_r(M,\mathcal{T}')} \delta_{c(e_1)c(e_3)} \prod_{e\in E'\setminus\{e_1,e_2\}} |e|_c \prod_{\Delta\in T'\setminus\{\Delta_1',\Delta_2'\}} |\Delta|_c = \sum_{c\in A_r(M,\mathcal{T})} \prod_{e\in E} |e|_c \prod_{\Delta\in T} |\Delta|_c,$$

which by definition equals $TV_r(M,\mathcal{T})$.

For the latter case, we suppose that $\mathcal{T}'$ is an ideal triangulation of M obtained from an ideal triangulation $\mathcal{T}$ by doing a $2-3$ Pachner move. Let E and T respectively be the sets of edges and tetrahedral of $\mathcal{T}$, and let E' and T' be the sets of edges and tetrahedra of $\mathcal{T}'$. Suppose $\Delta_1 \cup \Delta_2$ is where the $2-3$ move is performed, and $e_1 \in E'$ and Δ_1', Δ_2', $\Delta_3' \in T'$ are respectively the new edge and tetrahedral in $\mathcal{T}'$. Then $E' \setminus \{e_1\} = E$ and $T' \setminus \{\Delta_1', \Delta_2', \Delta_3'\} = T \setminus \{\Delta_1, \Delta_2\}$. See the figure below.



By the Biedenharn-Elliot identity (2.2), we have

$$\begin{aligned} TV_r(M,\mathcal{T}') &= \sum_{c\in A_r(M,\mathcal{T}')} \prod_{e\in E'} |e|_c \prod_{\Delta\in T'} |\Delta|_c \\ &= \sum_{c\in A_r(M,\mathcal{T}')} \prod_{e\in E'\setminus\{e_1\}} |e|_c \prod_{\Delta\in T'\setminus\{\Delta'_1,\Delta'_2,\Delta'_3\}} |\Delta|_c \Big(\sum_s |e_1|_s|\Delta'_1|_{c,s}|\Delta'_2|_{c,s}|\Delta'_3|_{c,s}\Big) \\ &= \sum_{c\in A_r(M,\mathcal{T}')} \prod_{e\in E'\setminus\{e_1\}} |e|_c \prod_{\Delta\in T'\setminus\{\Delta'_1,\Delta'_2,\Delta'_3\}} |\Delta|_c|\Delta_1|_c|\Delta_2|_c \\ &= \sum_{c\in A_r(M,\mathcal{T})} \prod_{e\in E} |e|_c \prod_{\Delta\in T} |\Delta|_c, \end{aligned}$$

which by definition equals $TV_r(M,\mathcal{T})$. □

4 Volume conjecture for 3-manifold with boundary

In this section, we provide numerical evidences for Conjecture 1.1. To simplify the notation, we will for each odd integer $r \geqslant 3$ let

$$QV_r(M) = \frac{2\pi}{r-2}\log\Big(TV_r(M; e^{\frac{2\pi\sqrt{-1}}{r}})\Big).$$

When M is the complement of a link L in S^3, we will let

$$TV_r(L) = TV_r(S^3\setminus L),$$

and let

$$QV_r(L) = QV_r(S^3\setminus L).$$

4.1 Figure-eight knot complement and its sister

The Figure-eight knot complement $S^3\setminus K_{4_1}$ has the following well known ideal triangulation,





from which Thurston [48] constructed the hyperbolic structure on $S^3\setminus K_{4_1}$, and calculated that

$$Vol(S^3\setminus K_{4_1}) \approx 2.02988.$$

In the figure above, edges with the same color (red or blue) are glued together and form an edge of the ideal triangulation. By Definition 3.1, for $r \geqslant 3$, we have

$$TV_r(K_{4_1}) = \sum_{(a,b)\in A_r} w_a w_b \begin{vmatrix} a & a & b \\ b & b & a \end{vmatrix}\begin{vmatrix} a & a & b \\ b & b & a \end{vmatrix},$$

where A_r consists of the pairs (a,b) of elements of I_r such that (a,a,b) and (b,b,a) are admissible, i.e., $2a-b\geqslant 0$, $2b-a\geqslant 0$, $2a+b\leqslant r-2$, $2b+a\leqslant r-2$ and $2a+b$ and $2b+a$ are integers. From this formula ,we have the following table of values of $QV_r(K_{4_1})$. Recall that we let $QV_r(M)=\frac{2\pi}{r-2}\log\big(TV_r(M;e^{\frac{2\pi\sqrt{-1}}{r}})\big)$.

r	11	13	15	17	19	21	23	25	31
$QV_r(K_{4_1})$	2.40661	2.37755	2.34826	2.31907	2.29953	2.28227	2.26834	2.25634	2.22824

r	41	51	61	71	81	91	101	111	121
$QV_r(K_{4_1})$	2.19685	2.17540	2.15953	2.14721	2.13731	2.12915	2.12230	2.11643	2.11136

r	131	141	151	201	301	401	501	701	1001
$QV_r(K_{4_1})$	2.10692	2.10299	2.09949	2.08641	2.07168	2.06344	2.05810	2.05153	2.04614

Figure 1 below is a comparison with the corresponding values of the Kashaev invariants $\langle K_{4_1}\rangle_r$, in which the red dots present the points $(r, QV_r(K_{4_1}))$, the blue dots represent the points $(r, \frac{2\pi}{r}\ln|\langle K_{4_1}\rangle_r|)$, the green dots represent the points $(r, Vol(S^3 \setminus K_{4_1}))$.



Figure 1: Comparison of different convergence modes for K_{4_1}

The manifold M_{2_2} in Callahan-Hildebrand-Weeks census [8], also known as the Figure-eight sister, shares the same volume with the Figure-eight knot complement, i.e.

$$Vol(M_{2_2}) = Vol(S^3 \setminus K_{4_1}) \approx 2.02988.$$

It is also known that M_{2_2} is not the complement of any knot in S^3. According to Regina [6], M_{2_2} has the following ideal triangulation.



Since for each tetrahedron in this triangulation, the coloring is the same as that of the one in the above ideal triangulation of $S^3 \setminus K_{4_1}$, $TV_r(M_{2_2})$ has exactly the same formula as $TV_r(K_{4_1})$, hence shares the same values.

4.2 Knot 5_2 complement and its sisters

According to SnapPy [16] and Regina [6], the knot 5_2 complement has volume

$$Vol(S^3 \setminus K_{5_2}) \approx 2.82812,$$

and has the following ideal triangulation.



Since only the colors of the edges (which edges are identified to form an edge in the triangulation) matters in the calculation of $TV_r(M)$, we omit the arrows on the edges. By Definition 3.1, for $r \geqslant 3$, we have

$$TV_r(K_{5_2}) = \sum_{(a,b,c)\in A_r} w_a w_b w_c \begin{vmatrix} a & a & b \\ b & c & c \end{vmatrix} \begin{vmatrix} a & a & b \\ b & c & c \end{vmatrix} \begin{vmatrix} a & b & c \\ b & b & c \end{vmatrix},$$

where A_r consists of triples (a,b,c) of elements of I_r such that (a,a,b), (b,b,c), (c,c,a) and (a,b,c) are admissible. From this ,we have the following table of values of $QV_r(K_{5_2})$.

r	7	9	11	21	31	41	51	61
$QV_r(K_{5_2})$	3.38531	3.32394	3.25282	3.09588	3.03657	3.00236	2.97925	2.96232
r	71	81	91	101	121	151	201	301
$QV_r(K_{5_2})$	2.94927	2.93883	2.93027	2.92309	2.91169	2.89937	2.88586	2.87071

Figure 2 below is a comparison with the corresponding values of the Kashaev invariants $\langle K_{5_2}\rangle_r$, in which the red dots present the points $(r, QV_r(K_{5_2}))$, the blue dots represent the points $(r, \frac{2\pi}{r}\ln|\langle K_{5_2}\rangle_r|)$, and the green dots represent the points $(r, Vol(S^3 \setminus K_{5_2}))$.

The manifold M_{3_6} in Callahan-Hildebrand-Weeks census [8], which is also known that M_{3_6} is the complement of the following knot with 12 crossings,



has the same volume as $S^3 \setminus K_{5_2}$, i.e.,

$$Vol(M_{3_6}) = Vol(S^3 \setminus K_{5_2}) \approx 2.82812.$$

According to Regina, M_{3_6} has an ideal triangulation as shown in the figure below. By the same reason, we still omit the arrows on the edges and keep the colors only.



Figure 2: Comparison of different convergence modes for K_{5_2}



Then for $r \geqslant 3$, we have

$$TV_r(M_{3_6}) = \sum_{(a,b,c)\in A_r} w_a w_b w_c \begin{vmatrix} a & a & b \\ b & c & c \end{vmatrix} \begin{vmatrix} a & a & b \\ b & c & c \end{vmatrix} \begin{vmatrix} a & b & c \\ a & a & c \end{vmatrix},$$

where A_r consists of triples (a,b,c) of elements of I_r such that (a,a,b), (b,b,c), (c,c,a), (a,b,c) and (a,a,c) are admissible. Below is a table of values $QV_r(K_{5_2})$ and $QV_r(M_{3_6})$, from which we see that the values of $QV_r(K_{5_2})$ and $QV_r(M_{3_6})$ are distinct, but are getting closer and closer as r grows, and both of them converge to 2.82812.

r	9	11	21	31	51	101
$QV_r(K_{5_2})$	3.3239396087031623282	3.2528240712684816477	3.0958786489268195966	3.0365668215995635907	2.9792536251826401549	2.9230944207585713174
$QV_r(M_{3_6})$	3.2936286562299185780	3.2291939333749922011	3.0954357480831343159	3.0365081953458580040	2.9792532229139281449	2.9230944207610719723

The manifold M_{3_8} in Callahan-Hildebrand-Weeks census [8] also has the same volume as $S^3 \setminus K_{5_2}$, and according to Regina has an ideal triangulation having the same colors as that of $S^3 \setminus K_{5_2}$. Therefore, $QV_r(M_{3_8})$ and $QV_r(K_{5_2})$ coincide for all $r \geqslant 3$. It is also known that M_{3_8} is not the complement of any knot in S^3.

4.3 Knot 6_1 complement

According to SnapPy [16] and Regina [6], the knot 6_1 complement has volume

$$Vol(S^3 \setminus K_{6_1}) \approx 3.163963,$$

and has the following ideal triangulation.



Since only the colors of the edges (which edges are identified to form an edge in the triangulation) matters in the calculation of $TV_r(M)$, we omit the arrows on the edges. By Definition 3.1, for $r \geqslant 3$, we have

$$TV_r(K_{6_1}) = \sum_{(a,b,c,d)\in A_r} w_a w_b w_c w_d \begin{vmatrix} a & a & b \\ a & d & b \end{vmatrix} \begin{vmatrix} a & c & c \\ b & b & d \end{vmatrix} \begin{vmatrix} b & b & c \\ a & c & d \end{vmatrix} \begin{vmatrix} b & b & c \\ b & d & c \end{vmatrix},$$

where A_r consists of triples (a, b, c, d) of elements of I_r such that all the triples involved are admissible. From this, we have the following table of values of $QV_r(K_{6_1})$.

r	5	7	9	11	21	31	41	51	61	71	81	91	101	121	151	201
$QV_r(K_{6_1})$	3.83348	3.63472	3.46573	3.39987	3.34732	3.31699	3.29688	3.28214	3.27076	3.26165	3.25417	3.24790	3.24255	3.23390	3.22431	3.21353

Figure 2 below is a comparison with the corresponding values of the Kashev invariants $\langle K_{6_1}\rangle_r$, in which the red dots present the points $(r, QV_r(K_{6_1}))$, the blue dots represent the points $(r, \frac{2\pi}{r}\ln|\langle K_{6_1}\rangle_r|)$, and the green dots represent the points $(r, Vol(S^3 \setminus K_{6_1}))$.

4.4 Some non-orientable cusped 3-manifolds

4.4.1 Gieseking manifold

Gieseking manifold is the smallest non-orientable cusped 3-manifold, which is the manifold N_{1_1} in Callahan-Hildebrand-Weeks census [8]. Therefore, it has an ideal triangulation with a single tetrahedron, and a Euler characteristic calculation shows that there is only one edge. According to SnapPy [16] and Regina [6], the Gieseking manifold has volume

$$Vol(N_{1_1}) \approx 1.014942.$$

By Definition 3.1, for $r \geqslant 3$, we have

$$TV_r(N_{1_1}) = \sum_{a\in A_r} w_a \begin{vmatrix} a & a & a \\ a & a & a \end{vmatrix},$$



Figure 3: Comparison of different convergence modes for K_{6_1}

where A_r consist of integers a such that $0 \leqslant a \leqslant (r-2)/3$. From this, we have the following table of values of $QV(N_{1_1})$.

r	7	9	11	21	31	41	51	61
$QV_r(N_{1_1})$	1.81736	1.66782	1.62276	1.43255	1.33012	1.27064	1.23174	1.20411
r	71	81	91	101	201	301	401	501
$QV_r(N_{1_1})$	1.18335	1.16711	1.15401	1.14319	1.08943	1.06872	1.05748	1.05035

4.4.2 Manifold N_{2_1}

According to SnapPy [16] and Regina [6], the manifold N_{2_1} in Callahan-Hildebrand-Weeks census [8] has volume

$$Vol(N_{2_1}) \approx 1.831931,$$

and has the following ideal triangulation.



By Definition 3.1, for $r \geqslant 3$, we have

$$TV_r(N_{2_1}) = \sum_{(a,b)\in A_r} w_a w_b \begin{vmatrix} a & b & b \\ a & b & b \end{vmatrix} \begin{vmatrix} a & b & b \\ a & b & b \end{vmatrix},$$

where A_r consist of the pairs $(a, b) \in I_r \times I_r$ such that (a, b, b) is addmissible. From this, we have the following table of values of $QV(N_{2_1})$.

r	5	7	9	11	21	31	41	51
$QV_r(N_{2_1})$	2.90345	2.54929	2.46119	2.42036	2.20099	2.11235	2.06163	2.02810
r	61	71	81	91	101	121	151	201
$QV_r(N_{2_1})$	2.00403	1.98578	1.97140	1.95974	1.95006	1.93489	1.91876	1.90140

4.5 Smallest hyperbolic 3-manifolds with totally geodesic boundary

By [34], each hyperbolic 3-manifold M_{min} with non-empty totally geodesic boundary that has the minimum volume admits a tetrahedral decomposition by two regular truncated hyperideal tetrahedra of dihedral angles $\pi/6$, and hence has volume

$$Vol(M_{\text{min}}) \approx 6.452.$$

Such minimum are not unique and are classified in [17]. In particular, the boundary of each M_{min} is a connected surface of genus 2, and an Euler number calculation shows that each ideal triangulation of M_{min} with two tetrahedra mush have only one edge. See the figure below.



Therefore, for $r \geqslant 3$,

$$TV_r(M) = \sum_{a \in A_r} w_a \begin{vmatrix} a & a & a \\ a & a & a \end{vmatrix} \begin{vmatrix} a & a & a \\ a & a & a \end{vmatrix},$$

where A_r consists of $a \in I_r$ such that (a, a, a) is admissible, i.e., a is an integer with $a \leqslant \lfloor (r-2)/3 \rfloor$. Here $\lfloor\ \rfloor$ is the floor function that $\lfloor x \rfloor$ equals the greatest integer that is less than or equal to x. Here for some r, it happens that $TV_r(M_{\text{min}})$ is negative. Since we require the argument to be in $[0, 2\pi)$ in the logarithm function, the imaginary part of $QV_r(M_{\text{min}})$ is always either 0 or $2\pi^2/(r-2)$, which converges to 0. Therefore, to test the convergence of $QV_r(M_{\text{min}})$, we can forget the imaginary parts and consider only the real parts. The table below contains some values of the real parts of $QV_r(M_{\text{min}})$.

r	11	21	31	41	51	61	71	81
$\Re(QV_r(M_{\text{min}}))$	4.39782	5.12434	5.44590	5.63235	5.75566	5.84395	5.91063	5.96297
r	91	101	201	301	401	501	1001	2001
$\Re(QV_r(M_{\text{min}}))$	6.00526	6.04022	6.21400	6.28075	6.31684	6.33970	6.38935	6.41741

Figure 4 below illustrates the asymptotic behavior of $QV_r(M_{\text{min}})$, where the red dots present the points $(r, QV_r(M_{\text{min}}))$ and the green dots represent the points $(r, Vol(M_{\text{min}}))$.



Figure 4: Asymptotics of $QV_r(M_{\text{min}})$

5 Volume conjecture for closed 3-manifolds

In this section, we provide numerical evidences for Conjectures 1.2 and 1.3 by calculating the Reshetikhin-Turaev-Lickorish invariants [45, 46, 31] of some oriented closed 3-manifolds. Recall that each oriented closed 3-manifold can be obtained from S^3 by doing an integral Dehn-surgery along a link. According to [31], if M is the 3-manifold obtained from S^3 by doing a p-surgery with $p \neq 0$ along a knot $K \subset S^3$, then for $r \geqslant 3$ odd, the Reshetikhin-Turaev-Lickorish invariants $\tau_r(M;q)$ of M at $q = e^{\frac{2\pi\sqrt{-1}}{r}}$ can be calculated by the following formula

$$\tau_r(M;q) = \frac{2}{r}e^{(\frac{3+r^2}{r}-\frac{3-r}{4})\pi\sqrt{-1}}\Bigg(\sum_{n=0}^{r-2}\big(\sin\frac{2(n+1)\pi}{r}\big)^2(-e^{\frac{\pi\sqrt{-1}}{r}})^{-p(n^2+2n)}J_{n+1}(K;e^{\frac{2\pi\sqrt{-1}}{r}})\Bigg), \qquad (5.1)$$

where $J_n(K;e^{\frac{2\pi\sqrt{-1}}{r}})$ is the value of the n-th colored Jones polynomial of K with 0-framing at $e^{\frac{2\pi\sqrt{-1}}{r}}$, normalized in the way that $J_n(\text{unknot}) = 1$.

Remark 5.1. Due to the convention difference of the quantum numbers $[n]$ we used here than that in some of the other literatures where $[n] = (q^{n/2} - q^{-n/2})/(q^{1/2} - q^{-1/2})$, the value $J_n(K;e^{\frac{2\pi\sqrt{-1}}{r}})$ we are going to calculate in the rest of the paper equals the value of the colored Jones polynomial at the root $e^{\frac{4\pi\sqrt{-1}}{r}}$ in their notation.

Remark 5.2. Formula (5.1) is derived directly from Lickorish ([31], §4.1 Definition). Using his notation and $A = e^{\frac{\pi\sqrt{-1}}{r}}$, one has $\mu = \frac{1}{\sqrt{r}}\sin\frac{2\pi}{r}$, $\langle\mu\omega\rangle^{\sigma}_{U_-} = e^{(\frac{3+r^2}{r}-\frac{3-r}{4})\pi\sqrt{-1}}$, $\langle\mu\omega\rangle^{-1}_{U} = \frac{2}{\sqrt{r}}\sin\frac{2\pi}{r}$ and for K_p the knot K with framing p,

$$\langle\omega\rangle_{K_p} = \sum_{n=0}^{r-2}\Bigg(\frac{\sin\frac{2(n+1)\pi}{r}}{\sin\frac{2\pi}{r}}\Bigg)^2(-e^{\frac{\pi\sqrt{-1}}{r}})^{-p(n^2+2n)}J_{n+1}(K;e^{\frac{2\pi\sqrt{-1}}{r}}).$$

Multiplying the above terms together, one gets formula (5.1).

To find the limit of the growth rates of $\tau_r(M;q)$, it is equivalent to calculate the limit as r approaches infinity of the following quantity

$$Q_r(M) = 2\pi\log\big(\tau_r(M;e^{\frac{2\pi\sqrt{-1}}{r}})/\tau_{r-2}(M;e^{\frac{2\pi\sqrt{-1}}{r-2}})\big).$$

5.1 Surgeries along Figure-eight knot

By (cf. [38]), the Figure-eight knot K_{4_1} has the n-th colored Jones polynomial

$$J_n(K_{4_1}, q) = \sum_{k=0}^{n-1} \prod_{i=1}^{k} (q^{n-i} - q^{-n+i})(q^{n+i} - q^{-n-i}). \tag{5.2}$$

In this subsection, we denote by M_p the manifold obtained from S^3 by doing a p-surgery along the K_{4_1}, and recall that M_p is hyperbolic when $|p| > 4$. In the tables below, we list $Q_r(M_p)$ for $p = -6, -5, 5, 6, 7$ and 8 and for $r = 51, 101, 151, 201, 301$ and 501.

5.1.1 p=-6

According to SnapPy [16],

$$Vol(M_{-6}) + CS(M_{-6})\sqrt{-1} = 1.28449 - 1.34092\sqrt{-1} \quad (mod \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.2), we have the following table of $Q_r(M_{-6})$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	101	151	201	301	501
$Q_r(M_{-6})$	$1.22717 - 1.34241\sqrt{-1}$	$1.28425 - 1.32879\sqrt{-1}$	$1.28440 - 1.33549\sqrt{-1}$	$1.28443 - 1.33786\sqrt{-1}$	$1.28446 - 1.33956\sqrt{-1}$	$1.28448 - 1.34043\sqrt{-1}$

5.1.2 p=-5

According to SnapPy,

$$Vol(M_{-5}) + CS(M_{-5})\sqrt{-1} = 0.98137 - 1.52067\sqrt{-1} \quad (mod \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.2), we have the following table of $Q_r(M_{-5})$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	101	151	201	301	501
$Q_r(M_{-5})$	$0.87410 - 1.50445\sqrt{-1}$	$0.98003 - 1.51521\sqrt{-1}$	$0.98130 - 1.51712\sqrt{-1}$	$0.98131 - 1.51865\sqrt{-1}$	$0.98134 - 1.51977\sqrt{-1}$	$0.98136 - 1.52035\sqrt{-1}$

5.1.3 p=5

According to SnapPy,

$$Vol(M_5) + CS(M_5)\sqrt{-1} = 0.98137 + 1.52067\sqrt{-1} \quad (mod \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.2), we have the following table of $Q_r(M_5)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	101	151	201	301	501
$Q_r(M_5)$	$0.87410 + 1.40967\sqrt{-1}$	$0.98003 + 1.49152\sqrt{-1}$	$0.98130 + 1.50659\sqrt{-1}$	$0.98131 + 1.51273\sqrt{-1}$	$0.98134 + 1.51714\sqrt{-1}$	$0.98136 + 1.51940\sqrt{-1}$

5.1.4 p=6

According to SnapPy,

$$Vol(M_6) + CS(M_6)\sqrt{-1} = 1.28449 + 1.34092\sqrt{-1} \quad (mod \ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.2), we have the following table of $Q_r(M_6)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	101	151
$Q_r(M_6)$	$1.22717 + 1.24762\sqrt{-1}$	$1.28425 + 1.30510\sqrt{-1}$	$1.28440 + 1.32496\sqrt{-1}$

r	201	301	501
$Q_r(M_6)$	$1.28443 + 1.33194\sqrt{-1}$	$1.28446 + 1.33693\sqrt{-1}$	$1.28448 + 1.33948\sqrt{-1}$

5.1.5 p=7

According to SnapPy,

$$Vol(M_7) + CS(M_7)\sqrt{-1} = 1.46378 + 1.19653\sqrt{-1} \quad (mod \ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.2), we have the following table of $Q_r(M_7)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	101	151
$Q_r(M_7)$	$1.43670 + 1.00605\sqrt{-1}$	$1.46354 + 1.15647\sqrt{-1}$	$1.46367 + 1.17877\sqrt{-1}$

r	201	301	501
$Q_r(M_7)$	$1.46372 + 1.18654\sqrt{-1}$	$1.46375 + 1.19209\sqrt{-1}$	$1.46377 + 1.19493\sqrt{-1}$

5.1.6 p=8

According to SnapPy,

$$Vol(M_8) + CS(M_8)\sqrt{-1} = 1.58317 + 1.07850\sqrt{-1} \quad (mod \ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.2), we have the following table of $Q_r(M_8)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	101	151
$Q_r(M_8)$	$1.57167 + 0.86833\sqrt{-1}$	$1.58282 + 1.03452\sqrt{-1}$	$1.58304 + 1.05896\sqrt{-1}$

r	201	301	501
$Q_r(M_8)$	$1.58309 + 1.06751\sqrt{-1}$	$1.58313 + 1.07362\sqrt{-1}$	$1.58315 + 1.07674\sqrt{-1}$

5.2 Surgeries along knot 5_2

By [35], the knot K_{5_2} has the n-th colored Jones polynomial

$$J_n(K_{5_2}, q) = \sum_{k=0}^{n-1} q^{\frac{k(k+3)}{2}} c_k \prod_{i=1}^{k} (q^{n-i} - q^{-n+i})(q^{n+i} - q^{-n-i}), \tag{5.3}$$

where

$$c_k = (-1)^k q^{\frac{5k^2+7k}{2}} \sum_{i=0}^{k} q^{i^2-2i-3ki} \frac{[k]!}{[i]![k-i]!}.$$

For simplicity, we denote by M_p the manifold obtained from S^3 by doing a p-surgery along the K_{5_2}, and recall that M_p is hyperbolic when $p \neq 0, 1, 2, 3$ or 4. In the tables below, we list $Q_r(M_p)$ for $p = -3, -2, -1, 5, 6$ and 7 and for $r = 51, 75, 101, 125, 151$ and 201.

5.2.1 p=-3

According to SnapPy [16],

$$Vol(M_{-3}) + CS(M_{-3})\sqrt{-1} = 2.10310 - 4.45132\sqrt{-1} \quad (mod\ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.3), we have the following table of $Q_r(M_{-3})$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	75	101	125	151	201
$Q_r(M_{-3})$	$2.10038 - 4.37951\sqrt{-1}$	$2.10200 - 4.41819\sqrt{-1}$	$2.10247 - 4.43323\sqrt{-1}$	$2.10268 - 4.43957\sqrt{-1}$	$2.10281 - 4.44329\sqrt{-1}$	$2.10293 - 4.44681\sqrt{-1}$

5.2.2 p=-2

According to SnapPy [16],

$$Vol(M_{-2}) + CS(M_{-2})\sqrt{-1} = 1.84359 - 4.63884\sqrt{-1} \quad (mod\ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.3), we have the following table of $Q_r(M_{-2})$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	75	101	125	151	201
$Q_r(M_{-2})$	$1.84822 - 4.59073\sqrt{-1}$	$1.84289 - 4.61357\sqrt{-1}$	$1.84317 - 4.62490\sqrt{-1}$	$1.84331 - 4.62978\sqrt{-1}$	$1.84339 - 4.63265\sqrt{-1}$	$1.84348 - 4.63536\sqrt{-1}$

5.2.3 p=-1

According to SnapPy [16],

$$Vol(M_{-1}) + CS(M_{-1})\sqrt{-1} = 1.39851 - 4.86783\sqrt{-1} \quad (mod\ \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.3), we have the following table of $Q_r(M_{-1})$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	75	101	125	151	201
$Q_r(M_{-1})$	$1.40943 - 4.84865\sqrt{-1}$	$1.39808 - 4.85045\sqrt{-1}$	$1.39817 - 4.85817\sqrt{-1}$	$1.39827 - 4.86157\sqrt{-1}$	$1.39834 - 4.86355\sqrt{-1}$	$1.39841 - 4.86542\sqrt{-1}$

5.2.4 p=5

According to SnapPy [16],

$$Vol(M_5) + CS(M_5)\sqrt{-1} = 0.98137 - 1.52067\sqrt{-1} \quad (mod \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.3), we have the following table of $Q_r(M_5)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	75	101	125	151	201
$Q_r(M_5)$	$0.87410 - 1.59924\sqrt{-1}$	$0.96890 - 1.53226\sqrt{-1}$	$0.98003 - 1.53890\sqrt{-1}$	$0.98098 - 1.53079\sqrt{-1}$	$0.98130 - 1.52765\sqrt{-1}$	$0.98131 - 1.52457\sqrt{-1}$

5.2.5 p=6

According to SnapPy [16],

$$Vol(M_6) + CS(M_6)\sqrt{-1} = 1.41406 - 1.51206\sqrt{-1} \quad (mod \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.3), we have the following table of $Q_r(M_6)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	75	101	125	151	201
$Q_r(M_6)$	$1.40044 - 1.56234\sqrt{-1}$	$1.41501 - 1.54958\sqrt{-1}$	$1.41339 - 1.53205\sqrt{-1}$	$1.41356 - 1.52508\sqrt{-1}$	$1.41372 - 1.52095\sqrt{-1}$	$1.41386 - 1.51706\sqrt{-1}$

5.2.6 p=7

According to SnapPy [16],

$$Vol(M_7) + CS(M_7)\sqrt{-1} = 1.75713 - 1.55255\sqrt{-1} \quad (mod \ \sqrt{-1}\pi^2\mathbb{Z}),$$

and by (5.1) and (5.3), we have the following table of $Q_r(M_7)$ modulo $\sqrt{-1}\pi^2\mathbb{Z}$.

r	51	75	101	125	151	201
$Q_r(M_7)$	$1.75178 - 1.63300\sqrt{-1}$	$1.75315 - 1.59624\sqrt{-1}$	$1.75507 - 1.57634\sqrt{-1}$	$1.75582 - 1.56798\sqrt{-1}$	$1.75625 - 1.56308\sqrt{-1}$	$1.75664 - 1.55846\sqrt{-1}$

6 Integrality conjecture for torus links

The purpose of this section is to provide supporting evidences to Conjecture 1.4, by both rigorous (§6.1) and numerical (§6.2) calculations.

6.1 Calculations for some torus links

In this subsection, we will calculate $TV_r(M)$ for the complement of some torus links, including the unknot, the Trefoil knot, the Hopf link and the torus links $T_{(2,4)}$ and $T_{(2,6)}$. To simplify the notation, for a link L in S^3, we let

$$TV_r(L) = TV_r(S^3 \setminus L).$$

All the ideal triangulations used in this section are obtain by using Regina [6] and SnapPy [16], and for simplicity, we will omit all the arrows on the edges and keep only the colors.

6.1.1 The unknot

Proposition 6.1. *Let U be the unknot in S^3. Then*

$$TV_r(U) = 1$$

for all $r \geqslant 3$ and for all roots of unity q of degree $2r$ such that q^2 is a primitive root of unity of degree r.

Proof. The unknot complement has the following ideal triangulation as shown in the figure below.



Therefore, for each $r \geqslant 3$, we have

$$TV_r(U) = \sum_{a,b} w_a w_b \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} \begin{vmatrix} a & a & a \\ a & a & a \end{vmatrix}, \tag{6.1}$$

where $(a, b) \in I_r \times I_r$ runs over all the admissible colorings at level r. The triple (a, a, a) being admissible implies that $a \in I_r \cap \mathbb{Z}$. Therefore, the right hand side of (6.1) equals

$$\sum_{a \in I_r \cap \mathbb{Z}} w_a \begin{vmatrix} a & a & a \\ a & a & a \end{vmatrix} \left(\sum_b w_b \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} \right),$$

where b is over all elements of I_r such that (a, b) is admissible at level r. Then the result follows from the following identity

$$\sum_b w_b \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} = \delta_{0,a}. \tag{6.2}$$

To prove (6.2), we use the Orthogonality. In (2.1), we let $m = 0$, $s = b$ and $i = j = k = l = n = a$. Then we have

$$\sum_b w_b w_0 \begin{vmatrix} a & a & 0 \\ a & a & b \end{vmatrix} \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} = \delta_{0,a},$$

where b is over all elements of I_r such that (a, b) is admissible at level r. Since $w_0 = 1$ and $\begin{vmatrix} a & a & 0 \\ a & a & b \end{vmatrix} = \frac{1}{[2a+1]}$, we have

$$\frac{1}{[2a+1]} \sum_b w_b \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} = \delta_{0,a}.$$

Since $a \leqslant (r-2)/2$, the number $1/[2a+1]$ is well defined and not equal to zero. As a consequence, (6.2) holds. □

Conjecture 6.2. *Let K be a knot in S^3. Then $TV_r(K) = 1$ for all r and q if and only if K is the unknot.*

Remark 6.3. It is interesting to find an $M \neq S^3 \setminus U$, not necessarily a knot complement, such that $TV_r(M) = 1$ for all r and q.

6.1.2 Trefoil knot

Proposition 6.4. *Let $T_{(2,3)}$ be the trefoil knot in S^3. Then*

$$TV_r(T_{(2,3)}) = \lfloor \frac{r-2}{3} \rfloor + 1,$$

for all $r \geqslant 3$ and for all roots of unity q of degree $2r$ such that q^2 is a primitive root of unity of degree r.

Proof. The $T_{(2,3)}$ knot complement has the following ideal triangulation as shown in the figure below.



Therefore, for each $r \geqslant 3$, we have

$$TV_r(T_{(2,3)}) = \sum_{a,b} w_a w_b \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix}, \tag{6.3}$$

where $(a, b) \in I_r \times I_r$ runs over all the admissible colorings at level r. The triple (a, a, a) being admissible implies that $a \in \mathbb{Z}$ and $a \leqslant (r-2)/3$. Hence the right hand side of (6.3) equals

$$\sum_{0 \leqslant a \leqslant \frac{r-2}{3}} \left(\sum_b w_b w_a \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} \right),$$

where a is over all the integers in that range and b is over all elements of I_r such that (a, b) is admissible at level r. Letting $i = j = k = l = m = n = a$ and $s = b$ in (2.1), we have

$$\sum_b w_b w_a \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} \begin{vmatrix} a & a & a \\ a & a & b \end{vmatrix} = 1,$$

where b is over all elements of I_r such that (a, a, b) is admissible at level r. As a consequence,

$$TV_r(T_{(2,3)}) = \sum_{0 \leqslant a \leqslant \frac{r-2}{3}} 1 = \lfloor \frac{r-2}{3} \rfloor + 1.$$

□

6.1.3 Hopf link and torus links $T_{(2,4)}$ and $T_{(2,6)}$

Proposition 6.5. *Let $T_{(2,2)}$ be the Hopf link in S^3. Then*

$$TV_r(T_{(2,2)}) = r - 1$$

for all $r \geqslant 3$ and for all roots of unity q of degree $2r$ such that q^2 is a primitive root of unity of degree r.

Proof. The Hopf link complement has the following ideal triangulation as shown in the figure below.







Therefore, for each $r \geqslant 3$, we have

$$
\begin{aligned}
TV_r(T_{(2,2)}) &= \sum_{a,b,c} w_a w_b w_c \begin{vmatrix} a & a & a \\ a & a & c \end{vmatrix} \begin{vmatrix} a & a & a \\ b & b & b \end{vmatrix} \begin{vmatrix} a & a & a \\ b & b & b \end{vmatrix} \\
&= \sum_{a,b} w_a w_b \begin{vmatrix} a & a & a \\ b & b & b \end{vmatrix} \begin{vmatrix} a & a & a \\ b & b & b \end{vmatrix} \left(\sum_{c} w_c \begin{vmatrix} a & a & a \\ a & a & c \end{vmatrix} \right),
\end{aligned}
$$

where for the first equation (a, b, c) runs over all the admissible colorings at level r, and in the second sum of the second equation c runs over all elements of I_r such that all the involved quantum $6j$-symbols are admissible at level r. By (6.2), we have

$$
\sum_{c} w_c \begin{vmatrix} a & a & a \\ a & a & c \end{vmatrix} = \delta_{0,a}.
$$

Therefore,

$$
TV_r(T_{(2,2)}) = \sum_{b} w_0 w_b \begin{vmatrix} 0 & 0 & 0 \\ b & b & b \end{vmatrix} \begin{vmatrix} 0 & 0 & 0 \\ b & b & b \end{vmatrix} = \sum_{b} 1,
$$

where the sum is over all elements b in I_r such that $(0, b, b)$ is admissible. Since $(0, b, b)$ is admissible for all $b \in I_r$, the sum is over all elements in I_r, and

$$
TV_r(T_{(2,2)}) = |I_r| = r - 1.
$$

□

Proposition 6.6. *Let $T_{(2,4)}$ be the $(2,4)$-torus link in S^3. Then*

$$
TV_r(T_{(2,4)}) = \Big(\lfloor \frac{r-2}{2} \rfloor + 1 \Big) \Big(\lfloor \frac{r-1}{2} \rfloor + 1 \Big)
$$

for all $r \geqslant 3$ and for all roots of unity q of degree $2r$ such that q^2 is a primitive root of unity of degree r.

Proof. The $T_{(2,4)}$ link complement has the following ideal triangulation as shown in the figure below.









Therefore, for each $r \geqslant 3$, we have

$$\begin{aligned} TV_r(T_{(2,4)}) &= \sum_{(a,b,c,d)\in A_r} w_a w_b w_c w_d \begin{vmatrix} a & a & b \\ c & c & c \end{vmatrix} \begin{vmatrix} a & a & b \\ c & c & c \end{vmatrix} \begin{vmatrix} b & b & b \\ a & a & d \end{vmatrix} \begin{vmatrix} b & b & b \\ a & a & a \end{vmatrix} \\ &= \sum_{a,b,c} w_a w_c \begin{vmatrix} a & a & b \\ c & c & c \end{vmatrix} \begin{vmatrix} a & a & b \\ c & c & c \end{vmatrix} \begin{vmatrix} b & b & b \\ a & a & a \end{vmatrix} \Big(\sum_d w_d w_b \begin{vmatrix} b & b & b \\ a & a & d \end{vmatrix} \Big), \end{aligned}$$

where in the second equation, a, b, c run over elements of I_r such that all the involved triples are admissible. We claim that

$$\sum_d w_d w_b \begin{vmatrix} b & b & b \\ a & a & d \end{vmatrix} = \sqrt{-1}^{2a+2b} \sqrt{[2a+1][2b+1]} \cdot \delta_{0,b}.$$

Indeed, letting $m = 0$, $i = j = a$, $k = l = n = b$ and $s = d$ in the Orthogonality (2.1), we have

$$\sum_d w_d w_b \begin{vmatrix} b & b & b \\ a & a & 0 \end{vmatrix} \begin{vmatrix} b & b & b \\ a & a & d \end{vmatrix} = \delta_{0,b}.$$

Then the claim follows from the fact that

$$\begin{vmatrix} b & b & b \\ a & a & 0 \end{vmatrix} = \frac{\sqrt{-1}^{2a+2b}}{\sqrt{[2a+1][2b+1]}}.$$

Therefore, we have

$$\begin{aligned} TV_r(T_{(2,4)}) &= \sum_{a,c} w_a w_c \begin{vmatrix} a & a & 0 \\ c & c & c \end{vmatrix} \begin{vmatrix} a & a & 0 \\ c & c & c \end{vmatrix} \begin{vmatrix} 0 & 0 & 0 \\ a & a & a \end{vmatrix} \sqrt{-1}^{2a} \sqrt{[2a+1]} \\ &= \sum_{a,c} (-1)^{2a}[2a+1](-1)^{2c}[2c+1] \frac{(-1)^{2a+2c}}{[2a+1][2c+1]} \frac{\sqrt{-1}^{2a}}{\sqrt{[2a+1]}} \sqrt{-1}^{2a} \sqrt{[2a+1]} \\ &= \sum_{a,c} 1, \end{aligned}$$

where a, c run over all the elements of I_r such that (c, c, a) and $(a, a, 0)$ are admissible. Counting the number of such pairs (a, c), we have

$$TV_r(T_{(2,4)}) = \Big(\lfloor \frac{r-2}{2} \rfloor + 1 \Big) \Big(\lfloor \frac{r-1}{2} \rfloor + 1 \Big).$$

□

Proposition 6.7. *Let $T_{(2,6)}$ be the $(2,6)$-torus link in S^3. Then*

$$TV_r(S^3 \setminus T(2,6)) = \Big(\lfloor \frac{r-2}{3} \rfloor + 1 \Big) \Big(\lfloor \frac{2r-2}{3} \rfloor + 1 \Big)$$

for all $r \geqslant 3$ and for all roots of unity q of degree $2r$ such that q^2 is a primitive root of unity of degree r.

Proof. The $T_{(2,6)}$ link complement has the following ideal triangulation as shown in the figure below.



Therefore, for each $r \geqslant 3$, we have

$$\begin{aligned} TV_r(T_{(2,6)}) =& \sum_{(a,b,c,d)\in A_r} w_a w_b w_c w_d \begin{vmatrix} a & a & c \\ b & b & a \end{vmatrix} \begin{vmatrix} a & a & c \\ b & b & a \end{vmatrix} \begin{vmatrix} b & b & c \\ b & b & b \end{vmatrix} \begin{vmatrix} b & b & d \\ b & b & c \end{vmatrix} \\ =& \sum_{a,b,c} w_a w_b \begin{vmatrix} a & a & c \\ b & b & a \end{vmatrix} \begin{vmatrix} a & a & c \\ b & b & a \end{vmatrix} \begin{vmatrix} b & b & c \\ b & b & b \end{vmatrix} \Big(\sum_d w_d w_c \begin{vmatrix} b & b & d \\ b & b & c \end{vmatrix} \Big), \end{aligned}$$

where in the second equation, a, b, c run over elements of I_r such that all the involved triples are admissible. We claim that

$$\sum_d w_d w_c \begin{vmatrix} b & b & d \\ b & b & c \end{vmatrix} = (-1)^{2b}[2b+1]\delta_{0,c}.$$

Indeed, letting $m = 0$, $i = j = k = l = b$, $n = c$ and $s = d$ in the Orthogonality (2.1), we have

$$\sum_d w_d w_c \begin{vmatrix} b & b & d \\ b & b & 0 \end{vmatrix} \begin{vmatrix} b & b & d \\ b & b & c \end{vmatrix} = \delta_{0,c}.$$

Then the claim follows from the fact that

$$\begin{vmatrix} b & b & d \\ b & b & 0 \end{vmatrix} = \frac{(-1)^{2b}}{[2b+1]}.$$

Therefore, we have

$$\begin{aligned} TV_r(T_{(2,6)}) =& \sum_{a,b} w_a w_b \begin{vmatrix} a & a & 0 \\ b & b & a \end{vmatrix} \begin{vmatrix} a & a & 0 \\ b & b & a \end{vmatrix} \begin{vmatrix} b & b & 0 \\ b & b & b \end{vmatrix} (-1)^{2b}[2b+1] \\ =& \sum_{a,b} (-1)^{2a}[2a+1](-1)^{2b}[2b+1] \frac{(-1)^{2a+2b}}{[2a+1][2b+1]} \frac{(-1)^{2b}}{[2b+1]} (-1)^{2b}[2b+1] \\ =& \sum_{a,b} 1, \end{aligned}$$

where a, b run over all the elements of I_r such that (a, a, b) and (b, b, b) are admissible. Counting the number of such pairs (a, b), we have

$$TV_r(T_{(2,6)}) = \Big(\lfloor \frac{r-2}{3} \rfloor + 1\Big)\Big(\lfloor \frac{2r-2}{3} \rfloor + 1\Big).$$

□

6.2 Numerical evidences for Conjecture 1.4

Apparently Propositions 6.1, 6.4, 6.5, 6.6, 6.7 are supporting evidences to Conjecture 1.4. In the rest of this subsection, we will provide more numerical evidences.

6.2.1 Knot $T_{(2,5)}$

The table below contains the value of $TV_r\big(T_{(2,5)}; e^{\frac{k\pi\sqrt{-1}}{r}}\big)$ for $k = 1, 2, 3$ and $r \leqslant 20$.

$k \setminus r$	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1	1	1	0.381966	1	3	3	2	0.763932	2	5	5	3	1.14590	3	7	7	4	1.52786
2	1		2.61803		3		2		2		5		7.85410		7		4	
3		1	2.61803		3	3		5.23607	2		5	3		3	7		4	10.4721

6.2.2 Knot $T_{(3,5)}$

The table below contains the value of $TV_r\left(T_{(3,5)}; e^{\frac{k\pi\sqrt{-1}}{r}}\right)$ for $k = 1, 2, 3$ and $r \leqslant 20$.

$k \setminus r$	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1	1	1	0.381966	1	2	3	2	1.38197	2	4	4	3	1.76393	3	6	6	4	2.14590
2	1		2.61803		2		2		2		4		6.23607		6		4	
3		1	2.61803		2	3		3.61803	2		4	3		3	6		4	8.85410

6.2.3 Knot $T_{(2,7)}$

The table below contains the value of $TV_r\left(T_{(2,7)}; e^{\frac{k\pi\sqrt{-1}}{r}}\right)$ for $k = 1, 2, 3$ and $r \leqslant 21$.

$k \setminus r$	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
1	1	1	2	1	0.307979	1	4	3	3	5	2	0.615957	2	7	5	5	8	3	0.923936
2	1		2		0.643104		4		3		2		2		5		8		1.92931
3		1	2		5.04892	1		3	3		2	10.0978		7	5		8	3	

6.2.4 Knot $T_{(3,7)}$

The table below contains the value of $TV_r\left(T_{(3,7)}; e^{\frac{k\pi\sqrt{-1}}{r}}\right)$ for $k = 1, 2, 3$ and $r \leqslant 21$.

$k \setminus r$	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
1	1	1	2	1	0.198062	1	3	3	3	4	2	0.841166	2	2	5	5	6	3	1.03923
2	1		2		3.24698		3		3		2		2		5		6		11.5429
3		1	2		1.55496	1		3	3		2	1.86294		2	5		6	3	

6.2.5 Knot $T_{(2,9)}$

The table below contains the value of $TV_r\left(T_{(2,9)}; e^{\frac{k\pi\sqrt{-1}}{r}}\right)$ for $k = 1, 2, 3$ and $r \leqslant 22$.

$k \setminus r$	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
1	1	1	1	2	3	1	0.283119	1	5	4	3	3	5	7	2	0.566237	2	9	7	5
2	1		1		3		0.426022		5		3		5		2		2		7	
3		1	1		3	1		1	5		3	3		7	2		2	9		5

6.2.6 Knot $T_{(2,11)}$

The table below contains the value of $TV_r\big(T_{(2,11)}; e^{\frac{k\pi\sqrt{-1}}{r}}\big)$ for $k = 1, 2, 3$ and $r \leqslant 22$.

$k \setminus r$	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
1	1	1	1	1	2	3	4	1	0.271554	1	6	5	4	3	3	5	7	9	2	0.543108
2	1		1		2		4		0.353253		6		4		3		7		2	
3		1	1		2	3		1	0.582964		6	5		3	3		7	9		1.16593

7 Some discussions and further questions

7.1 More comparisons with colored Jones polynomials

Let us take a closer look at the asymptotic behavior of $TV_r(M)$ in this subsection. To this end, we consider the following quantity

$$\Phi_r(M) \doteq \ln TV_r(M) - \frac{r-2}{2\pi} Vol(M).$$

For M being the K_{4_1} and K_{5_2} knot complements, by the data we obtained in §4.1 and §4.2, we found that the points $(\ln(r-2), \Phi_r(M))$ are asymptotic to a straight line $y = \frac{x}{2} + c$, where c is some constant. (See Figure 5 below, where in the first graph the points are respectively evaluated at $r = 101, 151, 201, 301, 401$ and 501, and in the second graph the points are respectively evaluated at $r = 81, 91, 101, 121, 151$ and 201.)



Figure 5

This suggests that

$$QV_r(K; e^{\frac{2\pi\sqrt{-1}}{r}}) = Vol(S^3 \setminus K) + \pi\frac{\ln(r-2)}{r-2} + O\Big(\frac{1}{r-2}\Big) \tag{7.1}$$

for K being the knots K_{4_1} or K_{5_2}. On the other hand, there is an expansion of colored Jones polynomial at roots of unity (cf. [56]),

$$J_r(K; e^{\frac{\pi\sqrt{-1}}{r}}) = 3^{-\frac{1}{4}} r^{\frac{3}{2}} e^{\frac{Vol(S^3\setminus K_{4_1})}{2\pi} r}\Big(1 + \frac{11\pi}{36\sqrt{3}r} + \cdots\Big),$$

which implies that

$$\frac{2\pi}{r}\ln|J_r(K;e^{\frac{\pi\sqrt{-1}}{r}})| = Vol(S^3\setminus K_{4_1}) + 3\pi\frac{\ln r}{r} + O(\frac{1}{r}). \tag{7.2}$$

Comparing (7.1) and (7.2), we see that in this case, $QV_r(K_{4_1};e^{\frac{2\pi\sqrt{-1}}{r}})$ converges much faster than $\frac{2\pi}{r}\ln|J_r(K;e^{\frac{\pi\sqrt{-1}}{r}})|$. It is really interesting to answer the following

Question 7.1. *Is the asymptotic behavior of $QV_r(K)$ for a hyperbolic knot always of the form*

$$QV_r(K;e^{\frac{2\pi\sqrt{-1}}{r}}) = Vol(S^3\setminus K) + \pi\frac{\ln(r-2)}{r-2} + O(\frac{1}{r-2})?$$

Or equivalently, is the asymptotic behavior of $TV_r(K)$ of the form

$$TV_r(K;e^{\frac{2\pi\sqrt{-1}}{r}}) = \alpha(r-2)^{1/2}e^{\frac{Vol(S^3\setminus K)}{2\pi}(r-2)}\Big(1+\frac{c_1}{r-2}+\frac{c_2}{(r-2)^2}+\cdots\Big)?$$

Proposition 6.1.2 and the data obtained in §6.2 suggest that $TV_r(K)$ grows linearly in $r-2$ for K being the torus knots $T_{(2,3)}$, $T_{(2,5)}$, $T_{(2,7)}$, $T_{(2,9)}$, $T_{(2,11)}$, $T_{(3,5)}$ and $T_{(3,7)}$, and Propositions 6.6 and 6.7 imply that $TV_r(T_{(2,4)})$ and $TV_r(T_{(2,6)})$ grow polynomially in $r-2$ of order 2. On the other hand, Kashaev-Tirkkonen [26] show that $|J_r(K;e^{\frac{2\pi\sqrt{-1}}{r}})|$ for a torus knot K grows in the order $r^{\frac{3}{2}}$, and Hikami [23] shows that $|J_r(L;e^{\frac{2\pi\sqrt{-1}}{r}})|$ for a torus link L of type $(2,2k)$ is deeply related to a class of modular forms of weight $3/2$. We are curious if there is some relationship between $TV_r(L)$ and certain modular forms. Combined with the cases of K_{4_1} and K_{5_2}, it shows that the asymptotic behavior of $TV_r(L)$ is quite different from that of the values of the colored Jones polynomial. Nonetheless, it is still very interesting to ask the following

Question 7.2. *Is there a relationship between $TV_r(S^3\setminus L)$ and the values of the colored Jones polynomials of L?*

As suggested by Proposition 6.1.2 and the data obtained in §6.2, it is interesting to answer the following

Question 7.3. *Let $T_{(m,n)}$ be a torus knot. Does $TV_r(T_{(m,n)})$ grow linearly in r?*

Recall that we consider the quantity

$$\Phi_r(M) = \ln TV_r(M) - \frac{r-2}{2\pi}Vol(M).$$

According to the data in §4.5, the points $(\ln(r-2), \Phi_r(M_{\min}))$ are asymptotic to a straight line $y = -\frac{3x}{2}+c$. (See the figure below, where the points are respectively evaluated at $r = 51, 101, 201, 301, 401, 501, 1001$ and 2001.)



This suggests that

$$QV_r(M_{\min};e^{\frac{2\pi\sqrt{-1}}{r}}) = Vol(M_{\min}) - 3\pi\frac{\ln(r-2)}{r-2} + O(\frac{1}{r-2}).$$

It is interesting to ask the following

Question 7.4. *Is the expansion above generally true for hyperbolic* 3*-manifolds* M *with totally geodesic boundary, or does the coefficient in front of* $\frac{\ln(r-2)}{r-2}$ *depend on the topology of* ∂M*?*

7.2 Other further questions

Question 7.5. *If* M *is obtained by gluing* M_1 *and* M_2 *together along some components of the boundary, do we have*

$$TV_r(M) = TV_r(M_1) \cdot TV_r(M_2)?$$

If so, then one could argue that the invariants $TV_r(M)$ *can be considered as the quantized version of the volume.*

Question 7.6. *Let* M *be a knot or link complement in* S^3*, how does* $TV_r(M)$ *behave under Dehn-fillings?*

Question 7.7. *For a compact* 3*-manifold* M *with boundary, what is the asymptotic behavior of* $\{TV_r(M; e^{\frac{\pi\sqrt{-1}}{r}})\}$*?*

Question 7.8. *For a compact* 3*-manifold* M *with or with out boundary, what is the asymptotic behavior of* $TV_r(M;q)$ *and, in the case of closed* M*, of* $\tau_r(M;q)$ *for the other roots of unity* q*? Numerical calculation suggests that at the other roots, the invariants also grow exponentially with the growth rate the volume.*

Question 7.9. *For a compact* 3*-manifold* M *with boundary, is there a Reshetikhin-Turaev type invariant* $\tau_r(M)$ *whose modulus gives* $TV_r(M)$*.*

Question 7.10. *Is there a definition of* $TV_r(S^3 \setminus L)$ *using a link diagram of* L*?*

Question 7.11. *Do there exist two distinct links* L_1 *and* L_2 *such that for all* $r \geqslant 3$*,*

$$TV_r(L_1) = TV_r(L_2)?$$

Question 7.12. *Is there a Turaev-Viro type invariant of closed* 3*-manifold defined using ideal triangulations and Thurston's spinning construction [48, 33]? If so, what is its relationship with the original Turaev-Viro invariant?*

References

[1] Baseilhac, Stphane; Benedetti, Riccardo, *Quantum hyperbolic invariants of* 3*-manifolds with* $PSL(2,\mathbb{C})$*-characters*, Topology 43 (2004), no. 6, 1373–1423.

[2] ———, *Classical and quantum dilogarithmic invariants of flat* $PSL(2,\mathbb{C})$*-bundles over* 3*-manifolds*, Geom. Topol. 9 (2005), 493–569.

[3] ———, *Quantum hyperbolic geometry*, Algebr. Geom. Topol. 7 (2007), 845–917.

[4] ———, *The Kashaev and quantum hyperbolic link invariants*, J. Gökova Geom. Topol. GGT 5 (2011), 31–85.

[5] Benedetti, Riccardo; Petronio, Carlo, *On Roberts' proof of the Turaev-Walker theorem*, J. Knot Theory Ramifications 5 (1996), no. 4, 427–439.

[6] Burton, Benjamin; Budney, Ryan; Pettersson, William, *Regina*, http://regina.sourceforge.net/.

[7] Callahan, Patrick J.; Dean, John C.; Weeks, Jeffrey R., *The simplest hyperbolic knots*, J. Knot Theory Ramifications 8 (1999), no. 3, 279–297.

[8] Callahan, Patrick J.; Hildebrand, Martin V.; Weeks, Jeffrey R., *A census of cusped hyperbolic* 3*-manifolds*, Math. Comp. 68 (1999), no. 225, 321–332.

[9] Casson, Andrew, private communications.

[10] Costantino, Francesco, *Coloured Jones invariants of links and the volume conjecture*, J. Lond. Math. Soc. (2) 76 (2007), no. 1, 1–15.

[11] ———, $6j$*-symbols, hyperbolic structures and the volume conjecture*, Geom. Topol. 11 (2007), 1831–1854.

[12] ———, *Integrality of Kauffman brackets of trivalent graphs,* Quantum Topol. 5 (2014), no. 2, 143–184.

[13] ———; Murakami, Jun, *On the $SL(2, \mathbb{C})$ quantum $6j$-symbols and their relation to the hyperbolic volume*, Quantum Topol. 4 (2013), no. 3, 303–351.

[14] ———; Guéritaud, François; van der Veen, Roland, *On the volume conjecture for polyhedra*, preprint: arXiv:1403.2347.

[15] ———; Geer, Nathan; Patureau-Mirand, Bertrand *Quantum invariants of 3-manifolds via link surgery presentations and non-semi-simple categories*, J Topology (2014) 7 (4): 1005–1053.

[16] Culler, Marc; Dunfield, Nathan, *SnapPy*, http:// sanppy.computop.org/.

[17] Fujii, Michihiko, *Hyperbolic 3-manifolds with totally geodesic boundary which are decomposed into hyperbolic truncated tetrahedra*, Tokyo J. Math. 13 (1990), no. 2, 353–373.

[18] Garoufalidis, Stavros; Lê, Thang T. Q., *On the volume conjecture for small angles*, preprint: arXiv:math/0502163.

[19] ———, *Asymptotics of the colored Jones function of a knot*, Geom. Topol. 15 (2011), no. 4, 2135–2180.

[20] Garoufalidis, Stavros; van der Veen, Roland, *Asymptotics of classical spin networks*, with an appendix by Don Zagier. Geom. Topol. 17 (2013), no. 1, 1–37.

[21] ———, *Asymptotics of quantum spin networks at a fixed root of unity*, Math. Ann. 352 (2012), no. 4, 987–1012.

[22] Gukov, Sergei, *Three-dimensional quantum gravity, Chern-Simons theory, and the A-polynomial*, Comm. Math. Phys. 255 (2005) 577–627.

[23] Hikami, Kazuhiro, *Quantum invariant for torus link and modular forms*, Comm. Math. Phys. 246 (2004), no. 2, 403–426.

[24] Kashaev, R. M., *A link invariant from quantum dilogarithm*, Modern Phys. Lett. A 10 (1995), no. 19, 1409–1418.

[25] ———, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. 39 (1997), no. 3, 269–275.

[26] ———; Tirkkonen, O., *A proof of the volume conjecture on torus knots*, (Russian) Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) 269 (2000), Vopr. Kvant. Teor. Polya i Stat. Fiz. 16, 262–268, 370; translation in J. Math. Sci. (N. Y.) 115 (2003), no. 1, 2033–2036

[27] Kauffman, Louis H.; Lins, Sstenes L., *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, Annals of Mathematics Studies, 134. Princeton University Press, Princeton, NJ, 1994.

[28] Kirillov, A. N.; Reshetikhin, N. Yu., *Representations of the algebra $U_q(sl(2))$, q-orthogonal polynomials and invariants of links*, Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988), 285–339, Adv. Ser. Math. Phys., 7, World Sci. Publ., Teaneck, NJ, 1989.

[29] Lickorish, W. B. R., *Three-manifolds and the Temperley-Lieb algebra*, Math. Ann. 290 (1991), no. 4, 657–670.

[30] ———, *Calculations with the Temperley-Lieb algebra. Comment. Math. Helv*, 67 (1992), no. 4, 571–591.

[31] ———, *The skein method for three-manifold invariants*, J. Knot Theory Ramifications 2 (1993), no. 2, 171–194.

[32] Luo, Feng, *A combinatorial curvature flow for compact 3-manifolds with boundary*. Electron. Res. Announc. Amer. Math. Soc. 11 (2005), 12–20 (electronic).

[33] ———; Tillmann, Stephan; Yang, Tian, *Thurston's spinning construction and solutions to the hyperbolic gluing equations for closed hyperbolic 3-manifolds*, Proc. Amer. Math. Soc. 141 (2013), no. 1, 335–350.

[34] Kojima, Sadayoshi; Miyamoto, Yosuke, *The smallest hyperbolic* 3*-manifolds with totally geodesic boundary*, J. Differential Geom. 34 (1991), no. 1, 175–192.

[35] Masbaum, Gregor, *Skein-theoretical derivation of some formulas of Habiro*, Algebr. Geom. Topol. 3 (2003), 537–556.

[36] Matveev, Sergei V., *Transformations of special spines, and the Zeeman conjecture*, (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 51 (1987), no. 5, 1104–1116, 1119; translation in Math. USSR-Izv. 31 (1988), no. 2, 423–434.

[37] ———, *Algorithmic topology and classification of* 3*-manifolds*, second ed., Algorithms and Com- putation in Mathematics, vol. 9, Springer, Berlin, 2007.

[38] Murakami, Hitoshi, *An introduction to the volume conjecture*, Interactions between hyperbolic geometry, quantum topology and number theory, 1–40, Contemp. Math., 541, Amer. Math. Soc., Providence, RI, 2011.

[39] ———; Murakami, Jun, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. 186 (2001), no. 1, 85–104.

[40] ———; Okamoto, Miyuki; Takata, Toshie; Yokota, Yoshiyuki, *Kashaev's conjecture and the Chern-Simons invariants of knots and links*, Experiment. Math. 11 (2002), no. 3, 427–435.

[41] ———; Yokota, Yoshiyuki, *The colored Jones polynomials of the figure-eight knot and the volumes of three-manifolds obtained by Dehn surgeries*, J. Reine Angew. Math. 607 (2007), 47–68.

[42] Murakami, Jun, *Generalized Kashaev invariants for knots in three manifolds*, preprint: arXiv:1312.0330.

[43] Ohtsuki, Tomotada, *Problems on invariants of knots and 3-manifolds*, with an introduction by J. Roberts. Geom. Topol. Monogr., 4, Invariants of knots and 3-manifolds (Kyoto, 2001), i–iv, 377–572, Geom. Topol. Publ., Coventry, 2002.

[44] Piergallini, Riccardo, *Standard moves for standard polyhedra and spines*, Third National Conference on Topology (Italian) (Trieste, 1986). Rend. Circ. Mat. Palermo (2) Suppl. No. 18 (1988), 391–414.

[45] Reshetikhin, N. Yu.; Turaev, V. G., *Ribbon graphs and their invariants derived from quantum groups*, Comm. Math. Phys. 127 (1990), no. 1, 1–26.

[46] ———, *Invariants of* 3*-manifolds via link polynomials and quantum groups*, Invent. Math. 103 (1991), no. 3, 547–597.

[47] Roberts, Justin *Skein theory and Turaev-Viro invariants*, Topology 34 (1995), no. 4, 771–787.

[48] Thurston, William, P., *Geometry and topology of* 3*-manifolds*, Revised Lecture Notes, Princeton Univ. Math. Dept. (1982)

[49] Turaev, Vladimir G., *Shadow links and face models of statistical mechanics*, J. Differential Geom. 36 (1992), no. 1, 35–74.

[50] ———, *Quantum invariants of links and* 3*-valent graphs in* 3*-manifolds*, Inst. Hautes tudes Sci. Publ. Math. No. 77 (1993), 121–171.

[51] ———, *Quantum invariants of knots and* 3*-manifolds*, Second revised edition. de Gruyter Studies in Mathematics, 18. Walter de Gruyter & Co., Berlin, 2010.

[52] ———; Viro, O. Ya., *State sum invariants of* 3*-manifolds and quantum* $6j$*-symbols*, Topology 31 (1992), no. 4, 865–902.

[53] Witten, Edward, *Quantum Field Theory and the Jones Polynomial*, Comm. Math. Phys. 121 (3): 351–399.

[54] van der Veen, Roland, *Proof of the volume conjecture for Whitehead chains*, Acta Math. Vietnam. 33 (2008), no. 3, 421–431.

[55] ———, *The volume conjecture for augmented knotted trivalent graphs*, Algebr. Geom. Topol. 9 (2009), no. 2, 691–722.

[56] Zagier, Don, *Quantum modular forms*, Quanta of Maths: Conference in honor of Alain Connes, Clay Mathematics Proceedings 11, AMS and Clay Mathematics Institute 2010, 659–675.

[57] Zheng, Hao, *Proof of the volume conjecture for Whitehead doubles of a family of torus knots*, Chin. Ann. Math. Ser. B 28 (2007), no. 4, 375–388.

Qingtao Chen
Mathematics Section, International Center for Theoretical Physics
Strada Costiera, 11, Trieste, I-34151, Italy
(qchen1@ictp.it)

Tian Yang
Department of Mathematics, Stanford University
Stanford, CA 94305, USA
(yangtian@math.stanford.edu)